

Radiative symmetry breaking and the $b \rightarrow s\gamma$ decay in generalized GMSB models

H. Hamidian^a, K. Huitu^b, K. Puolamäki^b, and D.-X. Zhang^b

^a*Department of Physics, Stockholm University, Box 6730, S-113 85 Stockholm, Sweden*

^b*Helsinki Institute of Physics, P.O.Box 9, FIN-00014 University of Helsinki, Finland*

Abstract

We study a class of generalized models of gauge mediated supersymmetry breaking (GMSB). We find the parameters and the full particle spectrum of the minimal supersymmetric standard model (MSSM) for all GMSB models with messenger multiplicities that satisfy the perturbativity of the gauge couplings up to the GUT scale. We give a detailed description of the algorithm that has been used to predict all the unmeasured parameters of the MSSM by taking (one-loop) radiative effects into account. We also calculate the branching ratio $BR(b \rightarrow s\gamma)$ and find that it always turns out to be larger than the standard model prediction. However, we find that the branching ratio typically remains within the current experimental bounds, except for some special cases with a light charged Higgs boson, or a small supersymmetry breaking scale and a negative μ -parameter.

11.30.Pb, 12.30.Qc, 14.80.Ly

A very important question in all supersymmetric (SUSY) grand unified theories beyond the standard model (SM) is to understand the mechanism of SUSY breaking and its communication to the observable (low-energy) sector. Although the answer to this question is still far from clear, there are a number of criteria which must be met regardless of any particular mechanism that breaks SUSY. For example, the so-called soft SUSY breaking mass terms must be small enough to ensure the stability of the Higgs mass against a large hierarchy of scales, while they must be large enough to have evaded the experimental searches so far.

SUSY breaking can be communicated to the visible sector either by gravitational interactions, as in supergravity (SUGRA) inspired models, or by SM gauge interactions, as in theories with gauge-mediated SUSY breaking (GMSB), which were initially studied nearly two decades ago [1]. The former models naturally lead to a universal soft-breaking sector near the Planck scale but are incapable of sufficiently suppressing large flavor violations at observable scales. Over the years many attempts have been made to solve this notorious SUSY flavor problem by invoking the “super-GIM” mechanism by assuming mass-degeneracy among the squarks and sleptons of a given flavor [2]. However, it is now understood that a number of rather generic phenomena, including non-minimal Kähler potentials, GUT effects, and superstring thresholds, can break degeneracy in SUGRA models and, in fact, even if all these could be somehow ruled out, scalar mass degeneracies will tend to be spoiled since flavor physics occurs below the scale at which SUSY-breaking is communicated to the SM. On the other hand, in GMSB theories gauge interactions provide flavor-symmetric SUSY breaking terms and thus naturally suppress the flavor-changing neutral currents associated with (universal) soft squark and slepton masses. Furthermore, due to the relatively low value of the SUSY-breaking scale in GMSB theories, this universality remains nearly unbroken as one evolves the mass parameters to the electroweak scale by using the renormalization group equations (RGEs). Among other attractive features of GMSB theories are the small number of free parameters, compared to the minimal supersymmetric standard model (MSSM), and the possibility of providing a solution to the SUSY CP problem. These features, together with the recent advances in understanding nonperturbative effects in SUSY gauge theories

[3] and the discovery of many new mechanisms for dynamical SUSY breaking (DSB) [4], have led to a revival of interest in GMSB models [5].

In the minimal version of the GMSB model [5] the messenger fields which communicate SUSY breaking to the visible sector belong to the $\mathbf{5} + \overline{\mathbf{5}}$ or $\mathbf{10} + \overline{\mathbf{10}}$ representations of the $SU(5)$ gauge group, and the messenger Yukawa couplings in any given $SU(5)$ representation are taken to be equal at the unification scale M_{GUT} . In order to communicate SUSY breaking, in addition to the particles in the minimal supersymmetric standard model (MSSM), the minimal GMSB theory contains at least one singlet superfield S which couples to vector-like messenger superfields $V + \overline{V}$ through the superpotential interaction

$$W_{\text{mess}} = \lambda_V S V \overline{V}. \quad (1)$$

At a scale $\Lambda_{\text{M}} \sim 10^2 - 10^4$ TeV, SUSY is broken and both the lowest and F -component, F_S , of the singlet superfield S acquire vacuum expectation values (VEVs) through their interactions with the hidden sector. The VEV $\langle S \rangle$ gives masses to the vector-like supermultiplets $V + \overline{V}$, while $\langle F_S \rangle$ induces mass splitting within the supermultiplets. Consequently, the gaugino and sfermion masses are generated through their gauge couplings to the messenger fields. The gauginos receive masses at one-loop, $m_\lambda \sim (\alpha/4\pi)\Lambda_{\text{SUSY}}$, where $\Lambda_{\text{SUSY}} = \langle F_S \rangle / \langle S \rangle$, while squarks and sleptons do so only at two-loop order, $\tilde{m}^2 \sim (\alpha/4\pi)^2 \Lambda_{\text{SUSY}}^2$. This implies that $m_\lambda \sim \tilde{m}$, which is another attractive feature of GMSB theories.

In the non-minimal generalizations of GMSB models [6] the messenger fields do not necessarily form complete $SU(5)$ GUT multiplets and one is naturally led to consider messenger fields which belong to incomplete representations of the $SU(5)$ gauge group. This can be seen by noting that the unification of the messenger Yukawa couplings at the GUT scale—whose MSSM analogue is the so-called $b - \tau$ unification [7]—is not necessarily required for gauge unification. For example, suppose that in addition to S there exist singlet superfields, S' , whose VEVs (but not the VEVs of their F -components) are just below the GUT scale, and which couple only to some components of the $SU(5)$ multiplet. Then, within the $SU(5)$ multiplet, these superfields acquire masses of order $\mathcal{O}(M_{\text{GUT}})$ and decouple from the low-

energy spectrum. The other components, which get their masses only through couplings with the superfield S , obtain masses of order $\lambda\langle S\rangle \sim \Lambda_{\text{M}}$. Since $\sqrt{\langle F_S\rangle}$ is much smaller than the masses of the heavy superfields, these (missing) particles make negligible mass contributions and play a less important role in determining the MSSM mass spectrum.

Although the phenomenological aspects of the minimal GMSB theories have been studied extensively [8–10], the generalized GMSB models—which may be more relevant for the construction of realistic SUSY GUTs—have been much less investigated. This is partly due to the larger messenger field content in the generalized GMSB theories which makes the study of these models technically more demanding. In a previous work [11] we have studied the phenomenological implications of generalized supersymmetric $SU(5)$ GUTs with GMSB by calculating the upper limits on $\tan\beta$ from nucleon decay and have found that the predicted values of $\tan\beta$ are mostly inconsistent with the constraints from nucleon decay. Our results suggest that in order to construct phenomenologically viable models, more complicated scenarios at the unification scale have to be considered.

In this Letter, we do not wish to speculate further on GUT scale physics in relation to GMSB theories, but instead take a bottom-up approach. The outline of our investigation is as follows: by assuming the most general sector of chiral messenger superfields, and restricting the number of messenger fields by demanding that the gauge couplings remain perturbative [6], 32 distinct possibilities for the messenger multiplicities are found. We assume vanishing bilinear and trilinear couplings in the soft scalar potential at the messenger scale. For each messenger sector there can be up to three different MSSM particle spectra and we list the full particle spectrum of all consistent models. In all the cases $\tan\beta$ is large and the μ -term can be either positive or negative, depending on the details of the messenger field spectrum. Our results are based on the calculation of the one-loop effective potential for a given messenger sector and we use the SM renormalization group equations (RGEs) to one-loop order to run the gauge and Yukawa couplings from the electroweak scale, m_Z , to the squark mass scale, m_{SUSY} .

We then compute the branching ratios for the $b \rightarrow s\gamma$ decay and compare it to the SM

prediction. This decay channel is very sensitive to new physics in the large $\tan\beta$ regime. The dominant non-SM contributions arise through constructive interference with the charged Higgs-boson loops and the next most important contributions are due to charginos. We find that in all the representative cases the branching ratios for the $b \rightarrow s\gamma$ decay are larger than the SM prediction. For the minimal GMSB models our results qualitatively agree with Borzumati's in [9].

Following Martin [6], we specify the messenger superfields in the generalized GMSB models as follows:

$$\begin{aligned}
n_Q : \quad Q + \overline{Q} &= (\mathbf{3}, \mathbf{2}, \frac{1}{6}) + \text{conj.}, \\
n_U : \quad U + \overline{U} &= (\overline{\mathbf{3}}, \mathbf{1}, -\frac{2}{3}) + \text{conj.}, \\
n_D : \quad D + \overline{D} &= (\overline{\mathbf{3}}, \mathbf{1}, \frac{1}{3}) + \text{conj.}, \\
n_L : \quad L + \overline{L} &= (\mathbf{1}, \mathbf{2}, -\frac{1}{2}) + \text{conj.}, \\
n_E : \quad E + \overline{E} &= (\mathbf{1}, \mathbf{1}, 1) + \text{conj.},
\end{aligned} \tag{2}$$

where the multiplicities of the messenger fields are denoted by $(n_Q, n_U, n_D, n_L, n_E)$.

At this stage, several restrictions can be used to reduce the number of possible models (see Ref. [6] for further discussion). By requiring that the gauge couplings remain perturbative, that the messenger field masses taking part in SUSY breaking not greatly exceed 10^4 TeV, and that all the gauginos acquire a non-vanishing mass, one ends up with 53 models obeying the following constraints on the messenger multiplicities [6]:

$$\begin{aligned}
(n_Q, n_U, n_D, n_L, n_E) &\leq (1, 0, 2, 1, 2) \\
&\text{or } (1, 1, 1, 1, 1) \\
&\text{or } (1, 2, 0, 1, 0) \\
&\text{or } (0, 0, 4, 4, 0).
\end{aligned} \tag{3}$$

Some of these 53 models give equivalent MSSM physics, since the sparticle masses are unaffected by changing $Q + \overline{Q} + E + \overline{E} \leftrightarrow 3(D + \overline{D} + L + \overline{L})$ (equivalent to $\mathbf{10} + \overline{\mathbf{10}} \leftrightarrow 3(\mathbf{5} + \overline{\mathbf{5}})$)

or $U + \bar{U} \leftrightarrow D + \bar{D} + E + \bar{E}$. Taking these permutation symmetries into account one is finally left with 32 different possibilities.

The vanishing trilinear and bilinear couplings in the soft scalar potential at the messenger scale makes GMSB models phenomenologically extremely predictive and fixes all the (as yet unmeasured) parameters of the MSSM for a given messenger sector. However, in order to implement this property one must compute the (radiatively corrected) effective potential and search for the minimum. We shall now proceed with the study of these radiative effects in the 32 models identified above and discuss their implications.

Our goal is to determine all the MSSM parameters for any given messenger sector. We specify the messenger sectors by their corresponding messenger multiplicities in (2) and by the messenger scale $\Lambda_M \simeq \lambda \langle S \rangle$. We have for simplicity taken only one effective messenger scale; all messengers have approximately the same mass. Generalization to multiple messenger scales would be straightforward.

For a given messenger sector, and at the messenger scale, we can calculate the bilinear term B (which is assumed to vanish) as a function of $\tan \beta = \langle H_u^0 \rangle / \langle H_d^0 \rangle$ and the sign of the μ -term. Thus, we must construct an algorithm for calculating the function $B(\tan \beta, \text{sign}(\mu))$ and then find its roots at Λ_M by standard numerical methods.

The one-loop effective potential written in terms of the VEVs, $v_u = \langle H_u^0 \rangle$ and $v_d = \langle H_d^0 \rangle$, is

$$V(Q) = V_0(Q) + \Delta V(Q), \quad (4)$$

where

$$\begin{aligned} V_0(Q) &= (m_{H_d}^2 + \mu^2) v_d^2 + (m_{H_u}^2 + \mu^2) v_u^2 - 2Bv_u v_d + \frac{1}{8} (g_L^2 + g_Y^2) (v_u^2 - v_d^2)^2, \\ \Delta V(Q) &= \frac{1}{64\pi^2} \sum_{k=\text{all the MSSM fields}} (-1)^{2S_k} n_k M_k^4 \left[-\frac{3}{2} + \ln \frac{M_k^2}{Q^2} \right], \end{aligned} \quad (5)$$

where S_k , n_k , and M_k are respectively the spin, the number of degrees of freedom, and the mass of the fields that contribute to ΔV , and where all the couplings are evaluated at some arbitrary renormalization scale Q .

The full effective potential is independent of the renormalization scale Q : the effects of changing the scale Q in $\Delta V(Q)$ are exactly cancelled as a result of the field and coupling constant renormalizations at the new scale. However, the one-loop leading-logarithmic approximation we are using is most accurate when the logarithmic terms in (5) are as small as possible, which is generally true when Q is chosen to equal $m_{\text{SUSY}} \simeq m_{\text{squark}}$.

First we use the SM RGEs to evolve the gauge and Yukawa couplings from the m_Z scale to the squark mass scale, m_{SUSY} . At the m_{SUSY} scale we match the SM and the MSSM parameters. This matching is straightforward at one-loop level, except for numerically significant two-loop corrections δ_b to the down quark Yukawa couplings in the high $\tan \beta$ regime [12]:

$$\begin{aligned} g_a^{\text{MSSM}}(m_{\text{SUSY}}) &= g_a^{\text{SM}}(m_{\text{SUSY}}), \\ Y_u^{\text{MSSM}}(m_{\text{SUSY}}) &= Y_u^{\text{SM}}(m_{\text{SUSY}}) / \sin \beta, \\ Y_{d33}^{\text{MSSM}}(m_{\text{SUSY}}) &= Y_{d33}^{\text{SM}}(m_{\text{SUSY}}) / ((1 + \delta_b) \cos \beta), \\ Y_{d,e}^{\text{MSSM}}(m_{\text{SUSY}}) &= Y_{d,e}^{\text{SM}}(m_{\text{SUSY}}) / \cos \beta. \end{aligned} \quad (6)$$

At first iteration we simply ignore the two-loop threshold effects and set $\delta_b = 0$. We use the MSSM RG equations to run the Yukawa couplings to the messenger scale Λ_{M} .

At the messenger scale the values of the mass-squared terms are given by the known two-loop graphs, while the trilinear soft terms vanish. We then run the Yukawa couplings, the mass-squared terms and the trilinear terms down to m_{SUSY} .

The minimization conditions for the tree-level potential V_0 are

$$\frac{\partial V_0}{\partial v_u} = \frac{\partial V_0}{\partial v_d} = 0. \quad (7)$$

Using (7) one can solve the bilinear term, which we denote by B_0 , and the μ -term, denoted by $\mu_0 = \text{sign}(\mu)\sqrt{\mu_0^2}$. As for δ_b , it can be approximated by [12]

$$\delta_b = \mu \tan \beta \left[\frac{2\alpha_3}{3\pi} M_3 I(m_{b_1}^2, m_{b_2}^2, M_3^2) + \frac{Y_{u33}}{16\pi^2} A_{u33} Y_{u33} I(m_{t_1}^2, m_{t_2}^2, \mu^2) \right], \quad (8)$$

where

$$I(x, y, z) = -\frac{xy \ln x/y + yz \ln y/z + zx \ln z/x}{(x-y)(y-z)(z-x)}. \quad (9)$$

At this stage we re-evaluate the down-Yukawa matrix at m_{SUSY} using (6) and (8), and repeat the previous steps to obtain the scalar mass-squared terms and the soft trilinear couplings at scale m_{SUSY} .

We then calculate the mass eigenvalues of all the gauge bosons, fermions, and scalars. We check that all the scalar mass-squared eigenvalues are non-negative, i.e., that we are at a true local minimum of the tree-level potential. The obtained masses and their first derivatives with respect to the VEVs are then used to minimize the full one-loop effective potential by using the equations

$$\frac{\partial V}{\partial v_u} = \frac{\partial V}{\partial v_d} = 0, \quad (10)$$

from which we can solve $\mu = \text{sign}(\mu)\sqrt{\mu_0^2 + \delta\mu^2}$ and $B = B_0 + \delta B$, where

$$\begin{aligned} \delta\mu^2 &= \frac{1}{2} \frac{v_d \partial \Delta V / \partial v_d - v_u \partial \Delta V / \partial v_u}{v_u^2 - v_d^2}, \\ \delta B &= \frac{1}{2} \frac{v_u \partial \Delta V / \partial v_d - v_d \partial \Delta V / \partial v_u}{v_u^2 - v_d^2}. \end{aligned} \quad (11)$$

Numerically, the most significant one-loop contribution to δB and $\delta\mu^2$ comes from the (s)top and (s)bottom loops. The contribution can be obtained by a straightforward calculation from eqs. (5) and (11) together with the quark masses $m_t = \lambda_t v_u$ and $m_b = \lambda_b v_d$ and the squark mass matrices.

Now all the parameters at the scale m_{SUSY} are known. We then finally run B and μ from m_{SUSY} to Λ_{M} . It remains to use numerical iteration to find the values of $\tan\beta$ and $\text{sign}(\mu)$ that satisfy $B_{\Lambda_{\text{M}}}(\tan\beta, \text{sign}(\mu)) = 0$, where we denote B at the scale Λ_{M} by $B_{\Lambda_{\text{M}}}$.

We use one-loop RGEs for all MSSM parameters, except for the bilinear scalar coupling B , for which we use the full two-loop expression [13]. Including the two-loop contribution to the running of B -parameter decreases (increases) $\tan\beta$ prediction for positive (negative) μ -parameter (for a discussion on including the two-loop RGE running see [9]).

We have listed all models consistent with radiative symmetry breaking in table I, taking $\Lambda_{\text{SUSY}} = 100 \text{ TeV}$ and $\Lambda_{\text{M}} = 100 \Lambda_{\text{SUSY}}$. There is a total of 32 possible messenger multiplicities and for each messenger multiplicity there can be several possible solutions corresponding

to different roots of the function $B_{\Lambda_M}(\tan\beta, \text{sign}(\mu))$.

In Fig. 1 we have plotted the function $B_{\Lambda_M}(\tan\beta, \text{sign}(\mu))$ for some representative models. In the minimal case $(0, 0, 1, 1, 0)$ there is only one root corresponding to a positive μ -term. There are three possible models having messenger multiplicities $(0, 0, 1, 2, 0)$, one root corresponding to positive μ -terms and two roots corresponding to a negative μ -term. For all messenger configurations the corresponding lines are cut above certain values of $\tan\beta$, since they would not lead to the desired radiative symmetry breaking: either the $SU(3)_C \times U(1)_{em}$ preserving vacuum state would actually be a saddle point of the effective potential due to the negative stau mass-squared eigenvalue, or the minimization conditions in eq. (10) would have no solutions for $|\mu^2| \geq 0$.

For every model we have calculated the ratio of the branching ratio $BR(b \rightarrow s\gamma)$ by using the full one-loop expression given in [14]. We find that the most important non-SM contribution comes from the constructive interference with the charged Higgs loop. For a positive (negative) μ -term the contribution from the chargino channel is typically approximately 5-30 % destructive (constructive). The gluino amplitude is typically about one per cent of the SM amplitude, while the neutralino contribution turns out to be totally negligible.

We find that in all the representative cases, except some models with very low SUSY breaking scale, the branching ratio $BR(b \rightarrow s\gamma)$ is larger than the SM prediction. The experimental bounds ($1.0 < 10^4 \times BR_{\text{EXP}}(b \rightarrow s\gamma) < 4.2$ [15]) combined with the theoretical uncertainty in the SM prediction ($10^4 \times BR_{\text{SM}}(b \rightarrow s\gamma) = 3.5 \pm 0.3$ [16]) limit the branching ratio to be between 0.3 and 1.4 times the SM prediction. At $\Lambda_{\text{SUSY}} = 100$ TeV, all models obey this limit.

In Fig. 2 we have plotted the branching ratios $BR(b \rightarrow s\gamma)$ for different messenger multiplicities and for different values of Λ_{SUSY} . The sparticle masses, which are shown in table I for $\Lambda_{\text{SUSY}} = 100$ TeV, are roughly proportional to the scale Λ_{SUSY} . For $\text{sign}(\mu) = +1$ the SM channel and the Higgs channel interfere destructively with the chargino channel. This interference keeps the branching ratio $BR(b \rightarrow s\gamma)$ approximately constant for all

scales for models with heavy charged Higgs fields. In the case $\text{sign}(\mu) = -1$ the chargino amplitude interferes constructively. As a result, the branching ratio can grow unacceptably large for small SUSY breaking scales.

To summarize, we have described in detail an algorithm to calculate all the unmeasured parameters of the MSSM for a given messenger sector with vanishing bilinear and trilinear soft terms. We have performed this calculation for a class of generalized GMSB models, giving the full particle spectrum for each model.

We have also calculated the branching ratios $BR(b \rightarrow s\gamma)$ for these models. We find that the branching ratio is almost always greater than the SM prediction. For most of these models the predicted branching ratios are less than 1.3 times the SM prediction. However, models with light ($m_H \lesssim 300$ GeV) charged Higgs boson, or small supersymmetry breaking scales and negative μ -parameter, result in unacceptably large $BR(b \rightarrow s\gamma)$.

ACKNOWLEDGMENTS

HH thanks the Swedish Natural Science Research Council for financial support. The work of KH, KP and DXZ is partially supported by the Academy of Finland (No. 37599).

We would like to thank F. Borzumati and T. Kobayashi for useful discussions and correspondence.

REFERENCES

- [1] M. Dine, W. Fischler, and M. Srednicki, Nucl. Phys. **B189** (1981) 575; S. Dimopoulos and S. Raby, Nucl. Phys. **B192** (1981) 353; M. Dine and W. Fischler, Phys. Lett. **110B** (1982) 227; M. Dine and M. Srednicki, Nucl. Phys. **B202** (1982) 238; L. Alvarez-Gaumé, M. Claudson, and M. Wise, Nucl. Phys. **B207** (1982) 96; C. Nappi and B. Ovrut, Phys. Lett. **113B** (1982) 175.
- [2] J. Donoghue, H.P. Nilles, and D. Wyler, Phys. Lett. **B128** (1983) 55; F. Gabbiani and A. Masiero, Nucl. Phys. **B322** (1989) 235; M. Dine, A. Kagan, and S. Samuel, Phys. Lett. **B243** (1990) 250; F. Gabbiani, E. Gabrielli, A. Masiero, and L. Silverstini, Nucl. Phys. **B477** (1996) 321.
- [3] N. Seiberg, Phys. Lett. **318B** (1993) 469; Phys. Rev. **D49** (1994) 6857; Nucl. Phys. **B435** (1995) 129; N. Seiberg and E. Witten, Nucl. Phys. **B426** (1994) 19, Erratum-*ibid.* **B430** (1994) 485; *ibid.* **B431** (1994) 484.
- [4] W. Skiba, Mod. Phys. Lett. **A12** (1997) 737.
- [5] M. Dine and A.E. Nelson, Phys. Rev. **D48** (1993) 1277; M. Dine, A.E. Nelson, and Y. Shirman, *ibid.* **D51** (1995) 1362; M. Dine, A.E. Nelson, Y. Nir, and Y. Shirman, *ibid.* **D53** (1996) 2658. See also, G.F. Giudice and R. Rattazzi, hep-ph/9801271 for review and further references.
- [6] S.P. Martin, Phys. Rev. **D56** (1997) 4207.
- [7] M. S. Chanowitz, J. Ellis, and M.K. Gaillard, Nucl. Phys. **B128** (1977) 506; R. Rattazzi, U. Sarid, and L.J. Hall, hep-ph/9405313.
- [8] S. Dimopoulos, M. Dine, S. Raby, and S. Thomas, Phys. Rev. Lett. **76** (1996) 3494; S. Dimopoulos, S. Thomas, and J. D. Wells, Phys. Rev. **D54** (1996) 3283; K. S. Babu, C. Kolda, and F. Wilczek, Phys. Rev. Lett. **77** (1996) 3070; J. A. Bagger, K. Matchev, D. M. Pierce, and R.-J. Zhang, Phys. Rev. **D55** (1997) 3188; H. Baer, M. Brhlik, C.-

- H. Chen, and X. Tata, Phys. Rev. **D55** (1997) 4463; N. G. Deshpande, B. Dutta, and S. Oh, Phys. Rev. **D56** (1997) 519.
- [9] R. Rattazzi and U. Sarid, Nucl. Phys. **B501** (1997) 297; F. M. Borzumati, hep-ph/9702307.
- [10] C. D. Carone and H. Murayama, Phys. Rev. **D53** (1996) 1658; B. Blok, C.-D. Lü, and D.-X. Zhang, Phys. Lett. **B386** (1996) 146.
- [11] H. Hamidian, K. Huitu, K. Puolamäki, and D.-X. Zhang, Phys. Lett. **B428** (1998) 310.
- [12] L. J. Hall, R. Rattazzi, and U. Sarid, Phys. Rev. **D50** (1994) 7048.
- [13] S. P. Martin and M. T. Vaughn, Phys. Rev. **D50** (1994) 2282.
- [14] S. Bertolini, F. Borzumati, A. Masiero, and G. Ridolfi, Nucl. Phys. **B353** (1991) 591.
- [15] M. S. Alam *et al.*, Phys. Rev. Lett. **74** (1995) 2885.
- [16] A.J. Buras, M. Misiak, M. Münz, and S. Pokorski, Nucl. Phys. **B424** (1994) 374.

TABLES

TABLE I. The MSSM particle spectrum for various messenger configurations ($\Lambda_{\text{SUSY}} = 100 \text{ TeV} = \Lambda_{\text{M}}/100$). The values of $\tan\beta$ and μ are calculated using one-loop effective potential, as described in text. The masses of the charged Higgs boson and spartners, expressed in GeV, are derived from the tree-level potential at the squark mass scale. At high $\tan\beta$ regime the lighter scalar Higgs boson mass is, after radiative corrections, about 130 GeV, while the pseudoscalar, charged and heavier scalar Higgs bosons are approximately mass-degenerate.

$(n_Q, n_U, n_D, n_L, n_E)$	$\tan\beta$	μ	$m_{H^\pm}$	$m_{\tilde{\chi}_{1,2}^\pm}$	$m_{\tilde{\chi}_{1,2,3,4}^0}$	$m_{\tilde{e}_{1,2}}$	$m_{\tilde{\tau}_{1,2}}$
	$\frac{\Gamma(b \rightarrow s\gamma)}{\Gamma_{SM}}$	M_3	$m_{\tilde{\nu}_e}/m_{\tilde{\nu}_\tau}$	$m_{\tilde{u}_{1,2}}$	$m_{\tilde{t}_{1,2}}$	$m_{\tilde{d}_{1,2}}$	$m_{\tilde{b}_{1,2}}$
(0, 0, 1, 1, 0)	36	504	511	248/503	136/248/491/502	192/362	141/370
	1.2	703	353/348	953/1006	821/945	947/1009	895/943
(0, 0, 1, 1, 1)	39	499	503	248/497	247/301/485/499	288/377	241/389
	1.2	703	369/362	963/1006	830/943	950/1010	892/941
(0, 0, 1, 1, 2)	41	493	495	248/492	247/447/479/512	365/394	309/414
	1.2	703	386/377	975/1007	839/941	953/1010	890/940
(0, 0, 1, 2, 0)	63	399	285	356/538	216/359/377/538	243/526	105/513
	1.3	698	520/500	954/1070	804/967	945/1073	820/958
(0, 0, 1, 3, 0)	27	-223	599	168/781	159/175/313/781	287/664	260/659
	1.4	695	660/654	956/1139	786/1066	943/1142	905/1055
(0, 0, 2, 1, 0)	21	788	804	253/775	193/253/769/774	226/364	198/375
	1.1	1364	356/354	1455/1488	1279/1414	1450/1490	1394/1438
(0, 0, 2, 1, 1)	22	785	802	253/771	253/359/765/771	313/380	282/397
	1.1	1364	372/369	1462/1488	1285/1413	1452/1490	1392/1438
(0, 0, 2, 1, 2)	23	781	801	253/767	253/524/761/769	387/397	343/427
	1.1	1364	389/386	1470/1489	1292/1413	1454/1491	1391/1439
(0, 0, 2, 2, 0)	38	724	726	499/716	276/499/701/716	272/525	218/528

	1.1	1359	519/511	1454/1530	1269/1433	1447/1532	1381/1431
(0, 0, 2, 3, 0)	55	644	569	596/790	358/598/615/790	313/662	202/651
	1.1	1355	657/639	1454/1577	1256/1449	1444/1579	1326/1443
(0, 0, 2, 4, 0)	66	537	297	494/1033	430/501/508/1033	351/787	160/761
	1.4	1352	783/753	1455/1628	1243/1469	1441/1630	1257/1460
(0, 0, 2, 4, 0)	23	-539	852	499/1033	432/505/511/1033	351/787	330/783
	1.2	1352	783/778	1455/1628	1244/1533	1441/1630	1408/1524
(0, 0, 3, 1, 0)	16	1015	1025	253/998	249/254/994/997	257/368	232/381
	1.1	2009	359/358	1902/1925	1691/1836	1897/1927	1819/1886
(0, 0, 3, 2, 0)	26	965	987	506/948	333/506/941/948	299/526	267/532
	1.1	2005	520/516	1900/1956	1683/1853	1893/1958	1832/1870
(0, 0, 3, 3, 0)	39	905	898	745/902	416/745/878/902	338/661	278/660
	1.1	2001	656/647	1899/1992	1674/1867	1890/1993	1811/1865
(0, 0, 3, 4, 0)	52	833	759	788/1038	499/790/802/1038	374/785	268/773
	1.1	1997	781/762	1898/2031	1663/1881	1887/2033	1762/1875
(0, 0, 4, 1, 0)	14	1216	1222	253/1199	253/306/1196/1198	285/372	259/388
	1.1	2645	363/362	2322/2339	2081/2236	2316/2341	2220/2307
(0, 0, 4, 2, 0)	21	1173	1199	507/1155	389/507/1150/1155	325/528	300/535
	1.1	2641	522/520	2320/2364	2075/2251	2313/2365	2235/2293
(0, 0, 4, 3, 0)	30	1123	1143	758/1108	473/758/1099/1108	362/662	322/664
	1.1	2637	657/651	2318/2392	2067/2264	2309/2394	2239/2275
(0, 0, 4, 4, 0)	41	1066	1047	973/1087	556/974/1039/1088	396/784	329/780
	1.1	2633	780/768	2317/2424	2058/2275	2306/2425	2213/2273
(1, 0, 0, 0, 1)	54	648	582	600/790	193/600/620/790	226/653	29/644
	1.1	1355	648/632	1447/1576	1250/1452	1442/1578	1329/1445
(1, 0, 0, 1, 0)	24	-548	832	509/1033	110/509/516/1033	171/772	122/768

	1.2	1352	768/763	1440/1627	1231/1531	1438/1629	1399/1522
(1, 0, 0, 1, 1)	23	-544	843	504/1033	275/506/511/1033	272/779	244/775
	1.2	1352	775/770	1447/1628	1237/1532	1439/1630	1404/1523
(1, 0, 0, 1, 1)	45	-542	478	501/1033	275/503/508/1033	272/779	173/767
	1.4	1352	775/762	1447/1628	1237/1482	1439/1630	1287/1474
(1, 0, 1, 0, 1)	38	908	905	745/905	250/745/882/904	257/652	184/651
	1.1	2001	647/638	1894/1991	1669/1869	1888/1993	1813/1867
(1, 0, 1, 1, 1)	51	837	770	792/1038	333/792/806/1038	299/777	167/766
	1.1	1997	773/756	1893/2031	1658/1882	1885/2032	1765/1877
(1, 0, 2, 0, 0)	29	1128	1152	758/1113	139/758/1104/1113	190/643	126/645
	1.1	2637	639/634	2309/2392	2059/2266	2307/2393	2242/2276
(1, 0, 2, 0, 1)	29	1126	1147	758/1111	306/758/1102/1110	285/652	241/654
	1.1	2637	647/642	2314/2392	2063/2265	2308/2393	2241/2276
(1, 0, 2, 1, 0)	39	1071	1062	976/1091	222/976/1045/1091	241/768	137/765
	1.1	2633	764/754	2307/2424	2050/2278	2304/2425	2217/2276
(1, 0, 2, 1, 1)	40	1069	1054	974/1089	390/975/1042/1089	325/776	248/772
	1.1	2633	772/761	2312/2424	2053/2277	2305/2425	2215/2274

FIGURES

FIG. 1. The values of the bilinear term B_{Λ_M} at the messenger scale as a function of $\tan\beta$. The thick lines correspond to the minimal model with messenger multiplicities $(0, 0, 1, 1, 0)$ at $\Lambda_{\text{SUSY}} = 100$ TeV and the thin lines correspond to the model with messenger multiplicities $(0, 0, 1, 2, 0)$ at $\Lambda_{\text{SUSY}} = 150$ TeV. The solid lines are for $\text{sign}(\mu) = +1$ and dashed lines are for $\text{sign}(\mu) = -1$. The minimal model has only one root consistent with the vanishing bilinear term at the messenger scale, while the other model has three, as can also be seen in figure 2. The lines are plotted up to the values of $\tan\beta$ below which the radiative symmetry breaking leads to the $SU(3)_C \times U(1)_{em}$ preserving physical vacuum state.




FIG. 2. The ratio $\Gamma(b \rightarrow s\gamma)/\Gamma_{SM}$ as a function of the scale Λ_{SUSY} ($\Lambda_M = 100 \Lambda_{SUSY}$). The solid lines correspond to the models with $\text{sign}(\mu) = +1$ and the dashed lines correspond to the models with $\text{sign}(\mu) = -1$. At $\Lambda_{SUSY} = 100$ TeV these lines correspond to the models listed in table I. Models with large rates are identified by $(n_Q, n_U, n_D, n_L, n_E)$ multiplicities.

